

MINIREVIEW

Amphiphilic Block Copolymers for Drug Delivery

MONICA L. ADAMS,¹ AFSANEH LAVASANIFAR,² GLEN S. KWON¹¹Division of Pharmaceutical Sciences, School of Pharmacy, University of Wisconsin–Madison, 777 Highland Avenue, Madison, Wisconsin 53705-2222²Faculty of Pharmacy and Pharmaceutical Sciences, University of Alberta, Edmonton, 3118 Dentistry/Pharmacy Centre, Alberta T6G 2N8, Canada*Received 18 November 2002; revised 16 December 2002; accepted 16 January 2003*

ABSTRACT: Amphiphilic block copolymers (ABCs) have been used extensively in pharmaceutical applications ranging from sustained-release technologies to gene delivery. The utility of ABCs for delivery of therapeutic agents results from their unique chemical composition, which is characterized by a hydrophilic block that is chemically tethered to a hydrophobic block. In aqueous solution, polymeric micelles are formed via the association of ABCs into nanoscopic core/shell structures at or above the critical micelle concentration. Upon micellization, the hydrophobic core regions serve as reservoirs for hydrophobic drugs, which may be loaded by chemical, physical, or electrostatic means, depending on the specific functionalities of the core-forming block and the solubilize. Although the Pluronics[®], composed of poly(ethylene oxide)-*block*-poly(propylene oxide)-*block*-poly(ethylene oxide), are the most widely studied ABC system, copolymers containing poly(L-amino acid) and poly(ester) hydrophobic blocks have also shown great promise in delivery applications. Because each ABC has unique advantages with respect to drug delivery, it may be possible to choose appropriate block copolymers for specific purposes, such as prolonging circulation time, introduction of targeting moieties, and modification of the drug-release profile. ABCs have been used for numerous pharmaceutical applications including drug solubilization/stabilization, alteration of the pharmacokinetic profile of encapsulated substances, and suppression of multidrug resistance. The purpose of this minireview is to provide a concise, yet detailed, introduction to the use of ABCs and polymeric micelles as delivery agents as well as to highlight current and past work in this area. © 2003 Wiley-Liss, Inc. and the American Pharmacists Association *J Pharm Sci* 92:1343–1355, 2003

Keywords: polymeric micelles; drug delivery; solubilization; PEG; polymer

INTRODUCTION

Many important therapeutic compounds exhibit poor aqueous solubility, rendering delivery of those agents quite challenging. The development

of effective delivery systems is crucial to the success of future drugs, which may include larger and more sophisticated synthetic compounds as well as complex natural molecules.¹

In addition to standard formulation techniques, polymeric micelles may be used for solubilization, stabilization, and delivery of challenging agents. The functional properties of micelles based on amphiphilic block copolymers (ABCs) render them ideal for encapsulation and delivery of

Correspondence to: Glen S. Kwon (Telephone: 608-265-5183; Fax: 608-262-5345; E-mail: gsk@pharmacy.wisc.edu)

Journal of Pharmaceutical Sciences, Vol. 92, 1343–1355 (2003)
© 2003 Wiley-Liss, Inc. and the American Pharmacists Association

hydrophobic drugs. ABCs consist of at least two regions of distinct chemical nature that undergo phase separation as a result of chain association in solvents that selectively dissolve one of the blocks. This process results in the formation of nanoscopic supramolecular core/shell structures, i.e., polymeric micelles. The assembly of ABCs into micelles can be largely explained by the relative interactions between the hydrophilic and hydrophobic blocks with one another and the surrounding medium. During the micellization process, the hydrophobic blocks associate to form the core region, whereas the hydrophilic segments position between the core and the external aqueous medium. Hence, the hydrophobic core is stabilized by the hydrophilic shell, which serves as an interface between the bulk aqueous phase and the hydrophobic domain (Fig. 1). This unique architecture enables polymeric micelles to serve as nanoscopic depots or stabilizers for poorly water-soluble compounds.

There has been great interest in the use of polymeric micelles as drug carriers.²⁻⁴ This mini-review will begin with an outline of the fundamental principles and basic characteristics of ABC delivery vehicles. The functional aspects of the hydrophilic shell-forming blocks and micellar dimensions will be outlined briefly. Emphasis will be given to three explicit categories of poly(ethylene oxide)-based ABCs, which are classified according to the chemical nature of the core-forming hydrophobic blocks. Therefore, the ABCs will be grouped as poly(ethylene oxide)-*block*-poly(L-amino acid)s, poly(ethylene oxide)-*block*-poly(ester)s, or poly(ethylene oxide)-*block*-poly(propylene oxide)-*block*-poly(ethylene oxide), i.e., the Pluronics®. Although the following discussion is focused primarily on self-associated delivery systems, other closely related work and the potential of ABC unimers will also be addressed. Past work, such as that involving the delivery and/or solubilization of challenging molecules, and more current efforts, such as those focused on gene delivery, will be highlighted.




Figure 1. Micellization model for an amphiphilic AB-diblock copolymer.

GENERAL CONSIDERATIONS AND PROPERTIES

Assembly

Polymeric micelles are association colloids. Hence, they are similar in some respects to standard surfactant micelles. Both ABCs and low-molecular-weight surfactants form micellar structures at or above a threshold critical micelle concentration (CMC). Below the CMC, the number of amphiphilic molecules adsorbing at the air-water interface increases with increasing concentration. At the CMC, both the bulk solution and the interface are saturated with unimers. For both types of amphiphiles, chain association is entropically driven via the expulsion of ordered water molecules into the bulk aqueous phase. The standard change in free energy for the micellization process, $\Delta G_{\text{mic}}^{\circ}$, is described by eq. 1:

$$\Delta G_{\text{mic}}^{\circ} = RT \ln(\text{CMC}) \quad (1)$$

where R represents the gas constant and T is the temperature of the system. The CMCs of ABCs are typically on the order of 10^6 – 10^7 M,^{5,6} whereas those of low-molecular-weight surfactants are on the order of 10^3 – 10^4 M.^{7,8} For this reason, micelles formed from ABCs are generally more thermodynamically stable than those formed from low-molecular-weight surfactants. Increased thermodynamic stability indicates that polymeric micelles should be less prone to disassembly at low concentrations than standard surfactants. ABC micelles also possess kinetic stability in that the dissociation of micelle structures into free-chain unimers is a slow process, even when the system is subjected to extreme dilution.^{9,10} Stability toward dilution is of particular concern for any association-based carrier system intended for parenteral administration and has implications for drug release. For example, a less thermodynamically and kinetically stable carrier might release drug prematurely because of dissociation of the micelle structure, whereas a more stable system might be preferred for long-circulating, sustained-release delivery systems.

Stealth Properties of Polymeric Micelles

A major barrier for colloidal drug carriers is non-specific uptake by the reticuloendothelial system (RES). The RES, or mononuclear phagocyte system, is a class of cells, including monocytes and

macrophages, responsible for engulfing and clearing old cells, miscellaneous cellular debris, foreign substances, and pathogens from the bloodstream. The ability to avoid RES uptake is crucial for achieving prolonged residence time in the blood compartment. For polymeric micelles, RES uptake is primarily a function of the hydrophilic shell, which is in direct contact with blood components after parenteral administration. The most commonly used hydrophilic block for polymeric micelle drug-delivery systems is poly(ethylene oxide)/poly(ethylene glycol), PEO/PEG. In the biomaterials literature, PEO is generally used in reference to polymers with a molecular weight $>10,000$ g/mol, whereas PEG is usually reserved for its lower molecular weight structural equivalent.¹¹⁻¹³ Presently, no distinction between the terms will be made and the authors' original nomenclature will be adhered to when referring to specific research.

PEO is FDA-approved for parenteral administration and is widely used in a variety of biomedical and pharmaceutical applications.^{14,15} One of the primary advantages of using PEO as a shell-forming material for polymeric micelles is low toxicity. For example, 10% PEG ($M_w = 4000$ g/mol) solutions have been safely administered intravenously to rats, guinea pigs, rabbits, and monkeys up to 16 g/kg.¹⁶ In addition, PEO has long been recognized for its ability to minimize protein adsorption to surfaces.^{11,13,17} Attachment of PEO chains to hydrophobic surfaces is particularly effective against protein adsorption because of the hydrophilicity and unique solution properties of PEO, including minimal interfacial free energy with water, high aqueous solubility, high mobility, and large exclusion volume.¹³ Therefore, PEO attachment is often used to improve the biocompatibility of foreign materials.

In PEO-based polymeric micelle systems, the attached PEO brushes impart steric stability by physically blocking interparticle attraction between core regions and, possibly, hindering interactions between the core-forming blocks and blood components. In particular, high surface density and long PEO chain length minimize protein adsorption to hydrophobic surfaces.¹⁷ For polymeric micelles, the length of the PEO blocks influences circulation time and uptake by the RES, with longer chains prolonging circulation time and decreasing RES uptake.¹⁸ Because PEO brushes are relatively inert and impart "stealth" properties to polymeric micelles, encapsulation in polymeric micelles may be a viable approach to

prolonging the circulation time of therapeutic agents.^{18,19}

Alternative Shell-Forming Materials

There has been some interest in developing other hydrophilic shell-forming polymers as PEO alternatives for polymeric micelles and liposomal systems.^{20,21} For example, Stepanek et al.²² have explored the use of mixed PEO/poly(electrolyte) block copolymers as shell-forming materials for polymeric micelles. In this system, it is possible to control micelle size via the relative expansion of the shell region, which is sensitive to changes in both pH and ionic strength. Furthermore, Roux et al.²³ recently demonstrated the effectiveness of a pH-sensitive *N*-isopropylacrylamide copolymer for steric stabilization of liposomes. The use of these types of hydrophilic polymers may have a more predominant role in polymeric micelle systems in the future. PEO alternatives may ultimately allow for tailoring of the shell properties, thereby increasing the utility of polymeric micelles for less traditional delivery applications, such as targeting, bioadhesion, responsiveness to proteins/cells, and pH sensitivity.

Micellar Dimensions

Any system intended for long circulation should be small enough so that it is not removed by filtration in the capillary beds.²⁴ Because polymeric micelles are usually <100 nm in diameter, they should be able to pass through microvasculature. The ability to avoid renal excretion may also have a role in prolonging blood residence time of polymeric micelles. Although the renal filtration limit has been explored most thoroughly for proteins and water-soluble polymers, the principles established in those realms are applicable to polymeric micelle systems. For proteins, there is some variation in the glomerular filtration limit because of charge, shape, etc. However, the cutoff is approximately 70,000 g/mol.²⁵ In most cases, the molecular weight of polymeric micelles is on the order of 10^6 g/mol.²⁶⁻²⁸ Therefore, characteristically high-molecular-weight polymeric micelles should avoid renal clearance. Conversely, ABC unimers, which fall below the glomerular filtration limit in terms of molecular weight, might be excreted upon micelle dissociation.

Generally, tumor vasculature is characterized by high permeability and decreased or defective

lymphatic drainage.²⁹ Consequently, it is possible for polymeric micelles, proteins, and other macromolecules to extravasate from leaky capillary beds *in vivo*, resulting in accumulation in pathological tissues that have poor lymphatic drainage.^{29,30} Depending on the tumor model, the endothelial pores of tumor vasculature vary in size from approximately 100 to 1000 nm.³¹ Yuan et al.³² demonstrated that tumor vasculature is six-fold more permeable to fluorescently labeled bovine serum albumin of approximately 5 nm than to sterically stabilized liposomes of 100 nm in diameter. Furthermore, the vasculature of human tumor xenografts implanted into immunocompromised mice is permeable to sterically stabilized liposomes of up to 400 nm.³³ Hence, the nanoscopic size of polymeric micelles may impart selectivity for tumor tissues, which should help minimize harmful side effects and toxicity. The ability of macromolecules to accumulate somewhat selectively in tumor tissues has been termed the enhanced permeation and retention (EPR) effect. The EPR effect has been exploited as a strategy for altering the biodistribution of encapsulated compounds by passively targeting polymeric micelles to solid tumors.¹⁸ An additional advantage associated with the small size of polymeric micelles is the relative ease of sterilization by filtration.

Core-Forming Blocks

In contrast to the ubiquitous use of PEO as a shell-forming material, numerous hydrophobic blocks have been studied for pharmaceutical applications. Thus, the core-forming blocks of polymeric micelle drug-delivery systems largely account for their flexibility and physicochemical properties. For clarity, the ABCs discussed in this minireview will be divided into three main categories based on the chemical structure of the core-forming block (Table 1). Specific ABC micelle systems may be further subclassified according to the method used to incorporate drugs or therapeutic biologicals. In general, the drug-loading method is determined primarily by the solubilities of the ABC and the solubilize. Commonly used loading methods include direct dissolution,³⁴ dialysis,^{35,36} oil/water emulsion,³⁷ chemical conjugation,³⁸ complexation,³⁹ and various solvent evaporation procedures.^{27,40} The most commonly used methods for micelle preparation using fairly water-soluble and water-insoluble ABCs are the direct dissolution and

dialysis methods, respectively. Depending on the method, drug loading may occur during or after micelle assembly. Furthermore, the manner in which drug is entrapped varies from system to system, depending on the method of loading and the ultimate goal of the research. Essentially, the mechanism of drug incorporation may be categorized as physical entrapment, chemical conjugation, or complexation. In some instances, polymer and drug may simply be administered congruently.⁴¹ The following discussion divides PEO-based polymeric micelle systems according to the chemical nature of the core-forming block. Specific properties and advantages of each class will be addressed.

POLY(ETHYLENE OXIDE)-BLOCK-POLY(L-AMINO ACID)S

Polymeric micelles based on poly(ethylene oxide)-*block*-poly(L-amino acid)s, PEO-*b*-p(L-AA)s, are quite versatile because they may facilitate chemical modification to the core-forming blocks and allow loading of therapeutic substances by both chemical and physical means. The hydrophobic portions of PEO-*b*-p(L-AA)s often contain functional groups that may be derivatized to enhance the properties of the core-forming blocks with respect to drug delivery. Consequently, the p(L-AA) block may offer the possibility of post-polymerization modification by chemical means.⁴²

Micelles based on poly(ethylene oxide)-*block*-poly(L-aspartate), PEO-*b*-p(L-Asp), have been used extensively for drug delivery. Because of the carboxyl functionality imparted by the p(L-Asp) block, PEO-*b*-p(L-Asp) is useful for the chemical conjugation of drugs.^{18,38} When drug is loaded in this manner, the polymer/drug conjugate acts as a prodrug in that cleavage of the drug by nonspecific hydrolysis or enzymatic activity is required for drug activity. This approach is appealing for the delivery of highly cytotoxic chemotherapeutic agents. In this case, polymer/drug conjugates are assembled into micelles, enabling drug molecules to be protected in the micelle core until the carrier vehicle accumulates at solid tumor sites due to the EPR effect.¹⁸ This strategy may minimize premature drug release and nonspecific action toward healthy cells, yet still allow for drug release in tumor tissues. Utilizing this approach with polymeric micelles may increase selectivity for tumor tissues via "passive" targeting.

Table 1. Block Copolymers Used for Drug and Gene Delivery

General Class	Incorporation Mechanism	Core-Forming Block	Encapsulated Compound	Reference
Poly(L-amino acid)	Chemical conjugation	Poly(L-aspartate)	Doxorubicin	18
		Poly(2-hydroxyethyl-L-aspartamide)	Methotrexate	38
	Physical entrapment	Poly(β -benzyl-L-aspartate)	Indomethacin	37
			KRN 5500	35
			Amphotericin B	44
	Complexation		Pyrene	45
			Doxorubicin	46
		Poly(<i>N</i> -hexyl-L-aspartamide)-acyl esters	Amphotericin B	27,28,40
		Poly(β -benzyl L-glutamate)	Clonazepam	43
		Poly(C ₁₆ -benzyl-L-aspartate)	KRN 5500	35
Poly(ester)	Chemical conjugation	Poly(L-aspartate-DOX)	Doxorubicin	47,48
		Poly(L-lysine)	Oligodeoxynucleotides	50
			DNA	52
		Poly(L-aspartate)	Cisplatin	39
			Lysozyme	51
	Physical entrapment	Poly(L-lysine)-succinate	Cisplatin	53
		Poly(D,L-lactic acid-co-glycolic acid)	Doxorubicin	69
		Poly(ϵ -caprolactone)	Indomethacin	6,58
			Dihydrotestosterone	36
			FK506	59
Pluronic [®]	Chemical conjugation		L-685,818	59
			Nimodipine	60
			Paclitaxel	63,64
			Doxorubicin	73
			Doxorubicin	69
	Physical entrapment	Poly(D,L-lactide)	Ruboxyl	82
		Poly(D,L-lactic acid-co-glycolic acid)	Doxorubicin	82,83
		n/a (P105)	Doxorubicin	34
		n/a (P85)	Doxorubicin	34,78,81
			Vinblastine	34
	Chemical conjugation		Mitomycin C	34
			Methotrexate	34
			Cisplatin	34
			Epirubicin	34
			Fluorescein	81
	Physical entrapment		Taxol	81
			Etoposide	81
			3'-azido-3'-deoxythymidin	81
			Valproic acid	81

(Continued)

Table 1. (Continued)

General Class	Incorporation Mechanism	Core-Forming Block	Encapsulated Compound	Reference
Cotreatment	n/a (F127) n/a (F68) n/a (P85)		Loperamide	81
			Nystatin	86
			Nystatin	86
			Digoxin	41
			Rhodamine-123	41,80
			Fluorescein	81
			Doxorubicin	81
			Taxol	81
			Etoposide	81
			3'-azido-3'-deoxythymidin	81
			Valproic acid	81
			Loperamide	81

Micelles based on PEO-*b*-p(L-Asp) have also been used for the physical entrapment of drugs. In this case, drug molecules may partition out of the micellar structure and assert pharmacological activity. Loading of amphotericin B (AmB), a potent antifungal agent, into PEO-*b*-p(*N*-hexyl-stearate-*L*-aspartamide) micelles results in reduced hemolytic activity compared with free AmB.⁴⁰ Furthermore, polymeric micelles based on PEO-*b*-poly(*N*-hexyl-*L*-aspartamide)-acyl esters solubilize AmB at high concentrations and result in sustained release of encapsulated drug *in vitro*.^{27,28} In one of the last examples, reconstitutable solid polymeric micelle formulations were prepared despite the relatively large size of the core-forming block.²⁷ The benzyl esters of both PEO-*b*-p(L-Asp) and PEO-*b*-(*L*-glutamate) also form micelles capable of encapsulating hydrophobic compounds and therapeutic agents.^{35,43-45} For instance, indomethacin, a poorly water-soluble drug, has been loaded into poly(ethylene oxide)-*block*-poly(β -benzyl-*L*-aspartate), PEO-*b*-PBLA.³⁷ It may be possible to use PEO-*b*-PBLA to take advantage of the EPR effect. To this end, Kataoka et al.⁴⁶ have loaded doxorubicin (DOX), a potent anticancer agent, into micelles prepared from PEO-*b*-PBLA.

Micelles prepared from derivatives of PEO-*b*-PBLA have also shown promise as delivery vehicles. For example, PEG-*b*-p(C₁₆-benzyl-*L*-aspartate) was prepared via an ester exchange reaction between PEO-*b*-PBLA and cetyl alcohol, and then used for incorporation of a water-insoluble anticancer drug.³⁵ To improve core/drug compatibility, it may even be possible to chemically tether the hydrophobic block of PEO-*b*-p(L-Asp) and other functional PEO-*b*-p(L-AA)s with the drug itself. To this end, Yokoyama et al.⁴⁷ derivatized PEO-*b*-p(L-Asp) via chemical conjugation of DOX and then loaded free DOX into micelles prepared from PEO-*b*-p(L-Asp-DOX). In this system, micelles containing high quantities of both chemically attached and physically loaded DOX express potent antitumor activity against a murine colon adenocarcinoma (C26) solid tumor model.⁴⁸ In a number of studies, the chemical structure of PEO-*b*-p(L-AA) has proven important in terms of drug-loading capacity, release characteristics, and drug activity. Thus, the ability to customize the core-forming block of functional PEO-*b*-p(L-AA)s is clearly advantageous.

Some PEO-*b*-p(L-AA)s also allow for ionic interactions between polymer chains. Therefore, it is

possible to prepare polyion complex (PIC) micelles, which contain charged polymer blocks, from block copolymers containing ionized p(L-AA) core-forming blocks. For instance, mixing oppositely charged polymers, such as negatively charged PEO-*b*-p(L-Asp) and positively charged PEO-*b*-p(L-Lysine), PEO-*b*-p(L-Lys), results in the formation of PIC micelles.⁴⁹ Similarly, it may be possible to take advantage of ionic interactions between therapeutic agents and polymers. For example, complexes formed between PEG-*b*-p(L-Lys) and antisense-oligodeoxynucleotides assemble into micellar structures with core/shell architecture.⁵⁰ PIC micelles may ultimately prove useful for antisense therapy or entrapment of enzymes.^{50,51}

Di- and triblock copolymers based on PEG-*b*-p(L-Lys) are currently being explored for gene delivery approaches focusing on nonviral transfection. Although micelles were not prepared, PEGylated linear, dendritic, and branched p(L-Lys) as well as PEG-grafted p(L-Lys) have been complexed with DNA and structure/activity relationships explored with respect to transfection efficiency.⁵² In this case, the physicochemical properties did not correlate with *in vitro* biological activity. However, such structure/property studies may aid researchers in developing block copolymer systems for use in gene delivery. Although delivery of DNA via complexation with PEG-*b*-p(L-Lys) has not yet been successful, exploitation of ionic interactions has shown promise as a means of drug loading. For example, metal complex formation can be utilized for loading cisplatin into PEO-*b*-p(L-Asp) micelles.³⁹ Cisplatin-loaded methoxy PEO-*b*-p(L-Lys)-succinate releases cisplatin slowly *in vitro* and alters the biodistribution of the drug.⁵³

An advantage of poly(L-AA)s is that they may be biodegradable and relatively nontoxic. The poly(amide) backbone of poly(L-AA)s may undergo hydrolysis and/or enzymatic degradation to yield biocompatible materials based on naturally occurring L-AAs. However, few toxicity/biocompatibility studies have been performed on PEO-*b*-p(L-AA)s and biodegradability remains to be established. Nonetheless, phase I clinical trials conducted in Japan suggest that DOX loaded into PEO-*b*-p(L-Asp) micelles containing a chemically conjugated DOX side chain is safe, with no overt signs of cardiotoxicity, liver toxicity, or untoward infusion-related side effects.⁵⁴ PEO-*b*-p(L-AA) systems have recently been reviewed in detail.⁵⁵

POLY(ETHYLENE OXIDE)- BLOCK-POLY(ESTER)S

ABCs containing poly(ester) core-forming blocks are often used for biomaterials and drug delivery applications. Although generally less suitable for chemical modification than poly(L-AA)s, poly(esters) have a history of safe application in humans. Aliphatic poly(ester)s are susceptible to nonspecific hydrolysis, and, therefore, subject to degradation in aqueous media.^{56,57} For delivery purposes, some of the most commonly used poly(ester)s include poly(glycolic acid), poly(D-lactic acid), poly(D,L-lactic acid), copolymers of lactide/glycolide, and poly(ϵ -caprolactone).

Although the core-forming block of PEO-*b*-p(ϵ -caprolactone), PEO-*b*-PCL, is semicrystalline, it is possible to control the degree of crystallinity of PCL by altering the ratio of PEO to PCL.⁵⁸ Allen et al.⁵⁹ have explored this system for formulation of immunosuppressive agents and demonstrated *in vitro* biocompatibility. In addition, dihydrotestosterone is effectively solubilized by PEO-*b*-PCL micelles and retains biological activity *in vitro*.³⁶ PEO-*b*-PCL micelles have also been used for encapsulation of indomethacin and result in sustained *in vitro* release.⁶ Ge et al.⁶⁰ demonstrated that triblock copolymers composed of PCL-*b*-PEO-*b*-PCL associate into micelle structures capable of encapsulating nimodipine, a lipophilic drug, in good yield.

Similar to micelles composed of PEO-*b*-p(L-AA)s, PEO-*b*-poly(ester) micelles may act as long-circulating drug carriers and allow for passive accumulation in tumor tissues via the EPR effect. Recently, the pharmacokinetic profile and biodistribution of PEO-*b*-PCL and PEO-*b*-PDLLA micelles have been investigated.^{61,62} Altering the pharmacokinetic profile of encapsulated agents is critical for exploiting the EPR effect as a means of targeting tumor tissues. PEO-*b*-poly(ester)s have been used extensively for solubilization of paclitaxel, a poorly water-soluble and potent anticancer agent. Methoxypoly(ethylene glycol)-*b*-poly(D,L-lactide), designated MePEG-*b*-PDLLA where Me represents the methoxy end-cap group of the PEG chain, is capable of solubilizing paclitaxel at >5000 times the saturation solubility of the drug in water.⁶³ Whereas MePEG-*b*-PDLLA micelles are nontoxic *in vivo*, MePEG-*b*-PDLLA unimers increase the apparent solubility of paclitaxel even below the CMC of the polymer.^{63,64} The basic and biological characteristics of poly(ether)-*b*-poly(ester) micelles for

formulation of paclitaxel have been reviewed by Liggins and Burt.⁶⁵

PEO-*b*-poly(ester)s have been used for other site-specific delivery applications. For example, PEG-*b*-PDLLA has recently been functionalized with aldehyde groups at the end of the PEG chains in order to introduce surface charge and amino acid residues to the micelle exterior.⁶⁶ In this way, it is possible to increase selectivity and further alter the biodistribution of polymeric micelles via the introduction of targeting moieties, such as sugars, to the periphery of the micelle shell.⁶⁷ With the PEG-*b*-PDLLA micelle system, Nagasaki et al.⁶⁸ conjugated various sugar molecules to the shell exterior and demonstrated selective protein interaction using affinity chromatography. The introduction of sugar moieties to the exteriors of polymeric micelle drug-delivery systems may ultimately allow for active targeting via glyco-receptors. Additionally, the thermal properties of PEG-*b*-PDLLA micelles have been exploited for drug delivery.²⁶

With PEO-*b*-poly(ester) micelle systems, it is possible to load drugs by both chemical and physical means. For example, PEG-*b*-poly(D,L-lactic acid-co-glycolic acid), PEG-*b*-PGLA, has been used for formulation of DOX through both drug conjugation and physical entrapment.⁶⁹ With this system, Yoo and Park⁶⁹ demonstrated different release profiles and postulated enhanced uptake of the drug conjugate. This system differs from those based on PEO-*b*-p(L-AA)s in that each polymer chain contains only one functional group at the end of the PGLA blocks for conjugation with DOX. Ultimately, this feature of PEG-*b*-PGLA may limit the attainable dose in animal models and humans.

In addition to their role as a core-forming material in traditional polymeric micelles, poly(esters) have often been incorporated into triblock copolymers for various delivery applications. In particular, poly(ester)s are often used in gel-forming systems. For example, triblock copolymers based on PEG-*b*-PGLA-*b*-PEG form biodegradable thermosensitive micelles.⁷⁰ With this system, it is possible to administer the formulation as a solution, which then gels at body temperature. Various biodegradable, poly(ester)-based gel-forming systems have been explored for delivery of therapeutic proteins.^{56,71} Recently, Kissel et al.⁷² have reviewed in detail the use of PEO-*b*-poly(ester)-*b*-PEO hydrogels for protein delivery. Similar triblock copolymers have been used for delivery of small molecules. For example, Liu et al.⁷³ have

prepared triblock copolymer micelles based on PDLLA-*b*-PEG-*b*-PDLLA for formulation of DOX and demonstrated sustained release characteristics *in vitro*.

POLY(ETHYLENE OXIDE)-BLOCK-POLY(PROPYLENE OXIDE)-BLOCK-POLY(ETHYLENE OXIDE)

Another important class of ABCs used in drug delivery applications is the Pluronics[®], also known as poloxamers. Pluronics[®] are triblock ABA-type copolymers based on poly(ethylene oxide)-*block*-poly(propylene oxide)-*block*-poly(ethylene oxide), which are typically expressed as PEO_{m/2}-*b*-PPO_n-*b*-PEO_{m/2}, where *m* and *n* designate the total average number of PEO and PPO repeat units, respectively. At low concentrations, Pluronics[®] alone are not cytotoxic.³⁴ A particularly attractive feature of the Pluronics[®] is that they are commercially available in a wide range of molecular weights and block ratios. The chemical structure of the copolymer, particularly the size of the PPO block, impacts both the CMC and partitioning of hydrophobic molecules into Pluronic[®] micelles.⁷⁴ Consequently, specific Pluronics[®] can be chosen so as to enhance drug-delivery properties.

Pluronics[®] have been shown to inhibit the efflux actions of P-glycoproteins (P-GP).^{34,75} The overexpression of P-GP in cancerous tissues is often linked to reduced accumulation of therapeutic agents at tumor sites.⁷⁵ P-GP-mediated multidrug resistance (MDR) is a mechanism through which tumors can become resistant to a variety of functionally and structurally diverse chemotherapies.⁷⁶ Pluronics[®] have shown potential for suppression of MDR. Specifically, the effect of Pluronics[®] on P-GP-mediated efflux is mediated primarily by the unimers.⁷⁷ The sensitization effect in resistant cells appears to be related to reduced adenosine 5'-triphosphate (ATP) levels in MDR phenotype cells, i.e., energy depletion.⁷⁸ Numerous cytotoxic agents, including both MDR- and non-MDR-type drugs, have been loaded into Pluronic[®] micelles (Table 1).^{34,77}

The use of Pluronics[®] may be a promising strategy for overcoming challenging drug delivery barriers, including MDR and the blood-brain barrier (BBB). The BBB is a barrier between the central nervous system (CNS) and systemic circulation, resulting from tight junctions between microvessel endothelial cells. Furthermore, efflux proteins decrease transcellular transport of

therapeutic compounds across the BBB. These characteristics make the BBB a substantial obstacle to CNS delivery.⁷⁹ Inhibition of the P-GP efflux transport system by Pluronics® may facilitate the delivery of therapeutic agents to the CNS.⁴¹ Pluronic® P85 induces ATP depletion and increased membrane fluidity in bovine microvessel endothelial cells, an *in vitro* model of the BBB.⁸⁰ In addition, the coadministration of digoxin and Pluronic® P85 increases the delivery of digoxin across the BBB *in vitro* by inhibiting P-GP-mediated drug transport and enhances delivery to the brain.⁴¹ Pluronic® P85, both above and below the CMC, has been used to increase transport of a number of therapeutic agents across a model BBB and Caco-2 cells.⁸¹

The use of ultrasonic enhancement has been explored as a method for controlled, or targeted, delivery with Pluronic® micelles. This approach makes use of the EPR effect to achieve selective accumulation at the target, i.e., tumor site. Then, ultrasound is applied exclusively to the tumor to yield localized drug delivery.⁸² Marin et al.⁸³ demonstrated that ultrasound enhances drug uptake *in vitro* by increasing both micelle uptake by HL-60 cells and the concentration of free drug in the incubation medium. This effect results in increased DOX uptake, even in MDR cells.⁸⁴ Like PEO-*b*-poly(ester)-*b*-PEO, Pluronics® have been used in gelling systems. For instance, mixed Pluronic® solutions containing F127 and F68 have been used for the development of a thermosetting ophthalmic drug-delivery device in order to increase ocular residence time.⁸⁵ Pluronics® F127 and F68 have also shown promise for formulation of nystatin, a polyene macrolide antifungal agent.⁸⁶ Using Pluronics®, it might be possible to attenuate the toxicity of nystatin via modulation of the self-aggregation state of the drug. Kabanov et al.^{87,88} have detailed the use of Pluronics® in both drug and gene delivery.

CONCLUSIONS

ABCs have proven useful and versatile for the delivery of therapeutic agents. In this regard, perhaps the most distinctive and functional aspect of ABCs is their ability to form micelles. Polymeric micelles are unique because of their characteristic nanoscopic core/shell architecture, and offer a viable means for the solubilization of hydrophobic compounds. Whereas the shell imparts steric stability and may possess targeting

moieties, the versatility and stability of ABC micelles results primarily from the chemical composition of the hydrophobic blocks. The chemical nature of the core-forming block may allow drug incorporation via chemical, physical, and/or electrostatic means. Depending on the application and desired release profile, more or less stable delivery systems might be necessary. Characteristically, micelles formed from ABCs are particularly stable to dilution and may exhibit prolonged circulation. Because diameters of <100 nm are generally considered prerequisite for avoiding RES uptake and increasing blood residence time, the size of polymeric micelles is a key factor in determining the biodistribution of encapsulated agents. Drugs may be encapsulated, and polymeric micelles may be prepared by a variety of techniques that include, but are not limited to, direct dissolution, dialysis, chemical complexation, O/W emulsion, and various solvent removal procedures. Once loaded, drugs are protected from the physiological environment, which might prohibit nonspecific and enzymatic hydrolysis, pH-dependent degradation, etc. Furthermore, drugs may be released in a controlled manner.

With all of these considerations in mind, it is possible to choose appropriate polymers and/or tailor ABCs to the task at hand, whether that be burst or sustained release, gene delivery, prolonged circulation, targeting, bioadhesion, etc. In some instances, unimers may be desirable and administered preferentially over micelles. In addition, it might be beneficial to reformulate existing compounds to minimize toxicity, alter the pharmacokinetic profile, and, ultimately, improve the therapeutic window of potent, yet toxic, compounds. Block copolymers are beginning to have a prominent role in gene delivery and might eventually be capable of replacing the viral vectors often used for such applications. To date, most research efforts have been focused on the hydrophobic core-forming blocks of ABCs. However, there is interest in the development of new shell-forming materials, which might elicit an appropriate response to a biological trigger, such as pH changes and protein recognition. Although exploitation of the EPR effect is often used for passive targeting, active targeting may have a more important role in future delivery efforts. Perhaps the introduction of targeting moieties to the hydrophilic shell-forming blocks will become more important and prevalent as researchers continue to strive for "smart" delivery systems that might be capable of targeting specific receptors or

cell types exclusively. The development of new polymers will surely lend greater flexibility and potential to delivery systems based on ABCs. The use of ABCs is a feasible and attractive alternative to standard formulation techniques. In particular, entrapment in polymeric micelles may provide a viable approach for stabilizing substances during systemic circulation. The use of ABCs for delivery purposes will likely increase as scientists are confronted with more and more challenging compounds and biologicals.

ACKNOWLEDGMENTS

The authors thank the National Institutes of Health for financial support (grant AI-43346-02). The authors also thank Karen Kostick of the University of Wisconsin-Madison School of Pharmacy, Division of Pharmaceutical Sciences, for her helpful suggestions.

REFERENCES

1. Davis SS, Illum L. 1998. Drug delivery systems for challenging molecules. *Int J Pharm* 176:1-8.
2. Jones M-C, Leroux J-C. 1999. Polymeric micelles: A new generation of colloidal drug carriers. *Eur J Pharm Biopharm* 48:101-111.
3. Rosler A, Vandermeulen GW, Klok H-A. 2001. Advanced drug delivery devices via self-assembly of amphiphilic block copolymers. *Adv Drug Deliv Rev* 53:95-108.
4. Kataoka K, Kwon G, Yokoyama M, Okano T, Sakurai Y. 1992. Block copolymer micelles as vehicles for drug delivery. *J Control Release* 24:119-132.
5. Kwon G, Naito M, Yokoyama M, Okano T, Sakurai Y, Kataoka K. 1993. Micelles based on ab block copolymers of poly(ethylene oxide) and poly(β -benzyl L-aspartate). *Langmuir* 9:945-949.
6. Kim SY, Shin IG, Lee YM, Cho CS, Sung YK. 1998. Methoxy poly(ethylene glycol) and ϵ -caprolactone amphiphilic block copolymeric micelles containing indomethacin. II. Micelle formation and drug release behaviours. *J Control Release* 51:13-22.
7. Myers D. 1999. Association colloids: Micelles, vesicles, and membranes. In: Myers D, editor. *Surfaces, interfaces, and colloids: Principles and applications*, 2nd ed. New York: John Wiley & Sons, p 383.
8. Goon P, Manohar C, Kumar VV. 1997. Determination of critical micelles concentration of anionic surfactants: Comparison of internal and external fluorescent probes. *J Colloid Interface Sci* 189:177-180.
9. Yokoyama M, Sugiyama T, Okano T, Sakurai Y, Naito M, Kataoka K. 1993. Analysis of micelle formation of an adriamycin-conjugated poly(ethylene glycol)-poly(aspartic acid) block copolymer by gel permeation chromatography. *Pharm Res* 10:895-899.
10. Halperin A, Alexander S. 1989. Polymeric micelles: Their relaxation kinetics. *Macromolecules* 22:2403-2412.
11. Wang P, Tan KL, Kang ET. 2000. Surface modification of poly(tetrafluoroethylene) films via grafting of poly(ethylene oxide) for reduction in protein adsorption. *J Biomater Sci Polym Ed* 11:169-186.
12. Zhang F, Kang ET, Neoh KG, Huang W. 2001. Modification of gold surface by grafting of poly(ethylene glycol) for reduction in protein adsorption and platelet adhesion. *J Biomater Sci Polym Ed* 12:515-531.
13. Lee JH, Lee HB, Andrade JD. 1995. Blood compatibility of polyethylene oxide surfaces. *Prog Polym Sci* 20:1043-1079.
14. Trubetskoy VS, Torchilin VP. 1995. Use of polyoxyethylene-lipid conjugates as long-circulating carriers for delivery of therapeutic and diagnostic agents. *Adv Drug Deliv Rev* 16:311-320.
15. Zalipsky S, Harris JM. 1997. Introduction to chemistry and biological applications of poly(ethylene glycol). In: Harris JM, Zalipsky S, editors. *Poly(ethylene glycol) chemistry and biological applications*, ACS symposium series 680. Washington, DC: American Chemical Society, pp 1-13.
16. Working PK, Newman MS, Johnson J, Cornacoff JB. 1997. Safety of poly(ethylene glycol) and poly(ethylene glycol) derivatives. In: Harris JM, Zalipsky S, editors. *Poly(ethylene glycol) chemistry and biological applications*, ACS symposium series 680. Washington, DC: American Chemical Society, pp 45-57.
17. Jeon SI, Lee JH, Andrade JD, De Gennes PG. 1991. Protein-surface interactions in the presence of polyethylene oxide. I. Simplified theory. *J Colloid Interface Sci* 142:149-158.
18. Kwon GS, Suwa S, Yokoyama M, Okano T, Sakurai Y, Kataoka K. 1994. Enhanced tumor accumulation and prolonged circulation times of micelle-forming poly(ethylene oxide-aspartate) block copolymer-adriamycin conjugates. *J Control Release* 29:17-23.
19. Yokoyama M, Okano T, Sakurai Y, Ekimoto H, Skibazaki C, Kataoka K. 1991. Toxicity and antitumor activity against solid tumors of micelle-forming polymeric anticancer drug and its extremely long circulation in blood. *Cancer Res* 51:3229-3236.
20. Taillefer J, Jones M-C, Brasseur N, Van Lier JE, Leroux J-C. 2000. Preparation and characterization

- of pH-responsive polymeric micelles for the delivery of photosensitizing anticancer drug. *J Pharm Sci* 89:52–62.
21. Torchilin VP, Levchenko TS, Whiteman KR, Yaroslavov AA, Tsatsakis AM, Rizos AK, Michailova EV, Shtilman MI. 2001. Amphiphilic poly-*N*-vinylpyrrolidones: Synthesis, properties, and liposome surface modification. *Biomaterials* 22: 3035–3044.
 22. Stepanek M, Podhajecka K, Tesarova E, Prochazka K. 2001. Hybrid polymeric micelles with hydrophobic cores and mixed polyelectrolyte/non-electrolyte shells in aqueous media. I. Preparation and basic characterization. *Langmuir* 17:4240–4244.
 23. Roux E, Stomp R, Giasson S, Pezolet M, Moreau P, Leroux J-C. 2002. Steric stabilization of liposomes by pH-responsive *N*-isopropylacrylamide copolymer. *J Pharm Sci* 91:1795–1802.
 24. Moghimi SM, Hunter AC, Murray JC. 2001. Long-circulating and target-specific nanoparticles: Theory to practice. *Pharmacol Rev* 53:283–318.
 25. Delgado C, Francis GE, Fisher D. 1992. The uses and properties of PEG-linked proteins. *Crit Rev Ther Drug Carrier Syst* 9:249–304.
 26. Yamamoto Y, Yasugi K, Harada A, Nagasaki Y, Kataoka K. 2002. Temperature-related change in the properties relevant to drug delivery of poly(ethylene glycol)-poly(D,L-lactide) block copolymer micelles in aqueous milieu. *J Control Release* 82: 359–371.
 27. Adams ML, Kwon GS. 2003. Relative aggregation state and hemolytic activity of amphotericin B encapsulated by poly(ethylene oxide)-*block*-poly(*N*-hexyl-L-aspartamide)-acyl conjugate micelles: Effects of acyl chain length. *J Control Release* 87:23–32.
 28. Lavasanifar A, Samuel J, Kwon GS. 2002. The effect of fatty acid substitution on the *in vitro* release of amphotericin B from micelles composed of poly(ethylene oxide)-*block*-poly(*N*-hexyl stearate-L-aspartamide). *J Control Release* 79:165–172.
 29. Maeda H, Wu J, Sawa T, Matsumura Y, Hori K. 2000. Tumor vascular permeability and the EPR effect in macromolecular therapeutics: A review. *J Control Release* 65:271–284.
 30. Maeda H, Sawa T, Konno T. 2001. Mechanism of tumor-targeted delivery of macromolecular drug, including the EPR effect in solid tumor and clinical overview of the prototype drug SMANCS. *J Control Release* 74:47–61.
 31. Hobbs SK, Monsky WL, Yuan F, Roberts WG, Griffith L, Torchilin VP, Jain RK. 1999. Regulation of transport pathways in tumor vessels: Role of tumor type and microenvironment. *Proc Natl Acad Sci* 95:4607–4612.
 32. Yuan F, Leunig M, Haung SK, Berk DA, Papahadjopoulos D, Jain RK. 1994. Microvascular permeability and interstitial penetration of sterically stabilized (stealth) liposomes in a human tumor xenograft. *Cancer Res* 54:3352–3356.
 33. Yuan F, Dellian M, Fukumura D, Leunig M, Berk DA, Torchilin VP, Jain RK. 1995. Vascular permeability in a human tumor xenograft: Molecular size dependence and cutoff size. *Cancer Res* 55:3752–3756.
 34. Alakhov VY, Moskaleva EY, Batrakova EV, Kabanov AV. 1996. Hypersensitization of multi-drug resistant human ovarian carcinoma cells by Pluronic P85 copolymer. *Bioconjug Chem* 7:209–216.
 35. Yokoyama M, Satoh A, Sakurai Y, Okano T, Matsumura Y, Kakizoe T, Kataoka K. 1998. Incorporation of water-insoluble anticancer drug into polymeric micelles and control of their particle size. *J Control Release* 55:219–229.
 36. Allen C, Han J, Yu Y, Maysinger D, Eisenberg A. 2000. Polycaprolactone-*b*-poly(ethylene oxide) copolymer micelles as a delivery vehicle for dihydrotestosterone. *J Control Release* 63:275–286.
 37. La SB, Okano T, Kataoka K. 1996. Preparation and characterization of the micelle-forming polymeric drug indomethacin-incorporated poly(ethylene oxide)-poly(β -benzyl L-aspartate) block copolymer micelles. *J Pharm Sci* 85:85–90.
 38. Li Y, Kwon GS. 2000. Methotrexate esters of poly(ethylene oxide)-*block*-poly(2-hydroxyethyl-L-aspartamide). I. Effects of the level of methotrexate conjugation on the stability of micelles and on drug release. *Pharm Res* 17:607–611.
 39. Nishiyama N, Kataoka K. 2001. Preparation and characterization of size-controlled polymeric micelle containing *cis*-dichlorodiammineplatinum (II) in the core. *J Control Release* 74:83–94.
 40. Lavasanifar A, Samuel J, Kwon GS. 2001. Micelles self-assembled from poly(ethylene oxide)-*block*-poly(*N*-hexyl stearate L-aspartamide) by a solvent evaporation method: Effect on the solubilization and haemolytic activity of amphotericin B. *J Control Release* 77:155–160.
 41. Batrakova EV, Miller DW, Li S, Alakhov VY, Kabanov AV, Elmquist WF. 2001. Pluronic P85 enhanced the delivery of digoxin to the brain: *In vitro* and *in vivo* studies. *J Pharmacol Exp Ther* 296:551–557.
 42. Adams ML, Kwon GS. 2002. The effects of acyl chain length on the micelle properties of poly(ethylene oxide)-*block*-poly(*N*-hexyl-L-aspartamide)-acyl conjugates. *J Biomater Sci Polym Ed* 13:991–1006.
 43. Jeong Y-I, Cheon J-B, Kim S-H, Nah J-W, Lee Y-M, Sung Y-K, Akaike T, Cho C-S. 1998. Clonazepam release from core-shell type nanoparticles *in vitro*. *J Control Release* 51:169–178.
 44. Yu BG, Okano T, Kataoka K, Sardari S, Kwon GS. 1998. *In vitro* dissociation of antifungal efficacy and toxicity for amphotericin B-loaded poly(ethylene

- oxide)-*block*-poly(β -benzyl-L-aspartate) micelles. *J Control Release* 56:285–291.
45. Kwon GS, Naito M, Kataoka K, Yokoyama M, Sakurai Y, Okano T. 1994. Block copolymer micelles as vehicles for hydrophobic drugs. *Colloids Surf B Biointerfaces* 2:429–434.
 46. Kataoka K, Matsumoto T, Yokoyama M, Okano T, Sakurai Y, Fukushima S, Okamoto K, Kwon GS. 2000. Doxorubicin-loaded poly(ethylene glycol)-poly(β -benzyl-L-aspartate) copolymer micelles: Their pharmaceutical characteristics and biological significance. *J Control Release* 64:143–153.
 47. Yokoyama M, Okano T, Sakurai Y, Kataoka K. 1994. Improved synthesis of adriamycin-conjugated poly(ethylene oxide)-poly(aspartic acid) block copolymer and formation of unimodal micellar structure with controlled amount of physically entrapped adriamycin. *J Control Release* 32:269–277.
 48. Yokoyama M, Fukushima S, Uehara R, Okamoto K, Kataoka K, Sakurai Y, Okano T. 1998. Characterization of physical entrapment and chemical conjugation of adriamycin in polymeric micelles and their design for *in vivo* delivery to a solid tumor. *J Control Release* 50:79–92.
 49. Harada A, Kataoka K. 1999. Chain length recognition: Core-shell supramolecular assembly from oppositely charged block copolymers. *Science* 283: 65–67.
 50. Harada A, Togawa H, Kataoka K. 2001. Physicochemical properties and nuclease resistance of antisense-oligodeoxynucleotides entrapped in the core of polyion complex micelles composed of poly(ethylene glycol)-poly(L-lysine) block copolymers. *Eur J Pharm Sci* 13:35–42.
 51. Harada A, Kataoka K. 2001. Pronounced activity of enzymes through the incorporation into the core polyion complex micelles made from charged block copolymers. *J Control Release* 72:85–91.
 52. Mannisto M, Vanderkerken S, Toncheva V, Elomaa M, Ruponen M, Schacht E, Urtti A. 2002. Structure-activity relationships of poly(L-lysines): Effects of PEGylation and molecular shape on physicochemical and biological properties in gene delivery. *J Control Release* 83:169–182.
 53. Bogdanov AA Jr, Martin C, Bogdanova AV, Brady TJ, Weissleder R. 1996. An adduct of *cis*-diamminedichloroplatinum (II) and poly(ethylene glycol)-poly(L-lysine)-succinate: Synthesis and cytotoxic properties. *Bioconjug Chem* 7:144–149.
 54. Matsumura Y. 2002. Phase I clinical trial of nk911, polymeric micelle encapsulated doxorubicin. In: Abstracts of the 2nd international symposium on tumor targeted delivery systems. Minneapolis, MN: Controlled Release Society, p 30.
 55. Lavasanifar A, Samuel J, Kwon GS. 2002. Poly(ethylene oxide)-*block*-poly(L-amino acid) micelles for drug delivery. *Adv Drug Deliv Rev* 54:169–190.
 56. Anseth KS, Metters AT, Bryant SJ, Martens PJ, Elisseeff JH, Bowman CN. 2002. *In situ* forming degradable networks and their application in tissue engineering and drug delivery. *J Control Release* 78:199–209.
 57. Wu XS, Wang N. 2001. Synthesis, characterization, biodegradation, and drug delivery application of biodegradable lactic-glycolic acid polymers. II. Biodegradation. *J Biomater Sci Polym Ed* 12: 21–24.
 58. Shin IG, Kim SY, Lee YM, Cho CS, Sung YK. 1998. Methoxy poly(ethylene glycol)/ ϵ -caprolactone amphiphilic block copolymeric micelles containing indomethacin. I. Preparation and characterization. *J Control Release* 51:1–11.
 59. Allen C, Yu Y, Maysinger D, Eisenberg A. 1998. Polycaprolactone-*b*-poly(ethylene oxide) block copolymer micelles as a novel drug delivery vehicle for neurotrophic agents FK506 and L-685,818. *Bioconjug Chem* 9:564–572.
 60. Ge H, Hu Y, Jiang X, Cheng D, Yuan Y, Bi H, Yang C. 2002. Preparation, characterization, and drug release behaviors of drug nimodipine-loaded poly(ϵ -caprolactone-poly(ethylene oxide)-poly(ϵ -caprolactone) amphiphilic triblock copolymer micelles. *J Pharm Sci* 91:1463–1473.
 61. Park YJ, Lee JY, Chang YS, Jeong JM, Chung JK, Lee MC, Park KB, Lee SJ. 2002. Radioisotope carrying polyethylene oxide-polycaprolactone copolymer micelles for targetable bone imaging. *Biomaterials* 23:873–879.
 62. Stolnik S, Heald CR, Neal J, Garnett MC, Davis SS, Illum L, Purkis SD, Barlow RJ, Gellert PR. 2001. Polylactide-poly(ethylene glycol) micellar-like particles as potential drug carriers: Production, colloidal properties, and biological performance. *J Drug Target* 9:361–378.
 63. Burt HM, Zhang X, Toleikis P, Embree L, Hunter WL. 1999. Development of copolymers of poly(D,L-lactide) and methoxypolyethylene glycol as micellar carriers of paclitaxel. *Colloids Surf B Biointerfaces* 16:161–171.
 64. Zhang X, Jackson JK, Burt HM. 1996. Development of copolymers as micellar carriers of Taxol. *Int J Pharm* 132:195–206.
 65. Liggins RT, Burt HM. 2002. Polyether-polyester diblock copolymers for the preparation of paclitaxel loaded polymeric micelle formulations. *Adv Drug Deliv Rev* 54:191–202.
 66. Yamamoto Y, Nagasaki Y, Kato M, Kataoka K. 1999. Surface charge modulation of poly(ethylene glycol)-poly(D,L-lactide) block copolymer micelles: Conjugation of charged peptides. *Colloid Surf B Biointerfaces* 16:135–146.
 67. Yasugi K, Nakamura T, Nagasaki Y, Kato M, Kataoka K. 1999. Sugar-installed polymer micelles: Synthesis and micellization of poly(ethylene glycol)-poly(D,L-lactide) block copolymers having sugar

- groups at the PEG chain end. *Macromolecules* 32:8024–8032.
68. Nagasaki Y, Yasugi K, Yamamoto Y, Harada A, Kataoka K. 2001. Sugar-installed block copolymer micelles: Their preparation and specific interaction with lectin molecules. *Biomacromolecules* 2:1067–1070.
 69. Yoo HS, Park TG. 2001. Biodegradable polymeric micelles composed of doxorubicin conjugate PLGA-PEG block copolymer. *J Control Release* 70: 63–70.
 70. Jeong B, Bae YH, Kim SW. 1999. Biodegradable thermosensitive micelles of PEG-PLGA-PEG triblock copolymers. *Colloid Surf B Biointerfaces* 16:185–193.
 71. Jeong B, Lee KM, Gutowska A, An YH. 2002. Thermogelling biodegradable copolymer aqueous solutions for injectable protein delivery and tissue engineering. *Biomacromolecules* 3:865–868.
 72. Kissel T, Li Y, Unger F. 2002. ABA-triblock copolymers from biodegradable polyester A-block and hydrophilic poly(ethylene oxide) B-blocks as a candidate for *in situ* forming hydrogel delivery systems for proteins. *Adv Drug Deliv Rev* 54:99–134.
 73. Liu L, Li C, Li X, Yuan Z, An Y, He B. 2001. Biodegradable polylactide/poly(ethylene glycol)/polylactide triblock copolymer micelles as anticancer drug carriers. *J Appl Polym Sci* 80:1976–1982.
 74. Kozlov MY, Melik-Nubarov NS, Batrakova EV, Kabanov AV. 2000. Relationship between Pluronic block copolymer structure, critical micellization concentration and partitioning coefficients of low molecular mass solutes. *Macromolecules* 33:3305–3313.
 75. Venne A, Li S, Madeville R, Kabanov A, Alakhov V. 1996. Hypersensitizing effect of Pluronic L61 on cytotoxic activity, transport, and subcellular distribution of doxorubicin in multiple drug-resistant cells. *Cancer Res* 56:3626–3629.
 76. Krishna R, Mayer LD. 2000. Multidrug resistance (MDR) in cancer. Mechanisms, reversal using modulators of MDR and the role of MDR modulators in influencing the pharmacokinetics of anticancer drugs. *Eur J Pharm Sci* 11:265–283.
 77. Batrakova E, Lee S, Li S, Venne A, Alakhov V, Kabanov A. 1999. Fundamental relationship between the composition of Pluronic block copolymers and their hypersensitization effect in MDR cancer cells. *Pharm Res* 16:1373–1379.
 78. Batrakova EV, Li S, Elmquist WF, Miller DW, Alakhov VY, Kabanov AV. 2001. Mechanism of sensitization of MDR cancer cells by Pluronic block copolymers: Selective energy depletion. *Br J Cancer* 85:1987–1997.
 79. Pardridge WM. 1995. Preface: Overview of brain drug delivery. *Adv Drug Deliv Rev* 15:1–3.
 80. Batrakova EV, Li S, Vinogradov SV, Alakhov VY, Miller DW, Kabanov AV. 2001. Mechanism of Pluronic effect on P-glycoprotein efflux system in blood-brain barrier: Contributions of energy depletion and membrane fluidization. *J Pharmacol Exp Ther* 299:483–493.
 81. Batrakova EV, Li S, Miller DW, Kabanov AV. 1999. Pluronic P85 increases permeability of a broad spectrum of drugs in polarized BBMEC and Caco-2 cell monolayers. *Pharm Res* 16:1366–1372.
 82. Rapaport NY, Herron JN, Pitt WG, Pitina L. 1999. Micellar delivery of doxorubicin and its paramagnetic analog, ruboxyl, to HL-60 cells: Effect of micelle structure and ultrasound on the intracellular drug uptake. *J Control Release* 58:153–162.
 83. Marin A, Muniruzzaman MD, Rapoport N. 2001. Acoustic activation of drug delivery from polymeric micelles: Effect of pulsed ultrasound. *J Control Release* 71:239–249.
 84. Marin A, Sun H, Hussein GA, Pitt WG, Christensen DA, Rapoport NY. 2002. Drug delivery in Pluronic micelles: Effect of high-frequency ultrasound on drug release from micelles and intracellular uptake. *J Control Release* 84:39–47.
 85. Wei G, Xu H, Ding PT, Li SM, Zheng JM. 2002. Thermosetting gels with modulated gelation temperature for ophthalmic use: The rheological and gamma scintigraphic studies. *J Control Release* 83:65–74.
 86. Yu B, Nichols BC, Aramwit P, Kwon G. 2000. The effect of Pluronics® F127 and F68 on the aggregation state and hemolytic activity of nystatin. In: Park KD, Kwon IC, Yui N, Jeong SY, Park K, editors. *Biomaterials and drug delivery toward new millennium*. Seoul: Han Rim Won Publishing, pp 373–381.
 87. Kabanov AV, Lemieux P, Vinogradov S, Alakhov V. 2002. Pluronic block copolymers: Novel functional molecules for gene therapy. *Adv Drug Deliv Rev* 54: 223–233.
 88. Kabanov AV, Batrakova EV, Alakhov VY. 2002. Pluronic block copolymers as novel polymer therapeutics for drug and gene delivery. *J Control Release* 82:189–212.